


Pvu II



5' ...CAG▼CTG...3'
3' ...GTC▲GAC...5'

PvuII is a restriction enzyme purified from a recombinant *E.coli* strain.

Catalogue No 128-1, 500 U
 128-2, 2500 U
 128-3, 12500 U

Concentration 10-12u/μl and 40-60u/μl*

*Add an H to cat.# to order the high concentration

Reagents supplied: 10x M and 10x K buffer

Unit substrate: Lambda DNA.

Unit calculation assay conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM dithiothreitol, 100 μg/ml bovine serum albumin and DNA. Incubate at 37°C.

Absence of contaminants: 100 units of *Pvu* II do not produce any unspecific cleavage products after 16 hrs incubation with 1 μg of Lambda DNA at 37°C. After 10-fold overdigestion with *Pvu* II greater than 95% of the DNA fragments can be ligated and recut with this enzyme.

Storage buffer: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 μg/ml bovine serum albumin and 50% glycerol. Store at -20°C.

Heat inactivation: No.

Methylation Sensitivity:

dam methylation: Not sensitive

dcm methylation: Not sensitive

CpG methylation: Not sensitive

Star activity: Conditions of low ionic strength, high enzyme concentration, glycerol concentration >5% or pH >8.0 may result in star activity.

Percent Activity in MINOTECH Buffers

L	M	H	SH	A	K
25-50	100	100	25-50	50	100

General reaction mixture:

10U PvuII 1μl
10x M or K buffer * 2μl
DNA substrate <1μg
Sterile ultrapure water Up to 20 μl
Incubate for 15 min at 37°C

*In the case of M buffer we recommend the addition of BSA to a final concentration of 100 μg/ml.

Frequency of Cutting

λ	Ad-2	Φx174	pUC18	M13mp18	pBR322
15	24	0	2	3	1



Lambda DNA 1.0 % agarose