


NruI



5' ...TCG▼CGA...3'
3' ...AGC▲GCT...5'

NruI is a restriction enzyme purified from *Nocardia rubra*.

Catalogue No 125-1, 1000 U
 125-2, 3x1000 U

Concentration 10-12u/μl and 40-60u/μl*

*Add an H to cat.# to order the high concentration

Reagents supplied: 10x U_{NruI} and 10x K buffer.

Unit substrate: Lambda DNA.

Unit calculation assay conditions: 100 mM KCl, 50 mM Tris-HCl (pH 8.0 @ 25°C), 10 mM MgCl₂, 100 μg/ml bovine serum albumin and DNA. Incubate at 37°C.

Absence of contaminants: 80 units of *Nru* I do not produce any unspecific cleavage products after 16 hrs incubation with 1 μg of Lambda DNA at 37°C. After 10-fold overdigestion with *Nru* I, less than 20% of the DNA fragments can be ligated.

Storage buffer: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 μg/ml bovine serum albumin and 50% glycerol. Store at -20°C.

Heat inactivation: 65°C for 20 minutes.

Methylation Sensitivity:

dam methylation: Blocked by overlapping
dcm methylation: Not sensitive
CpG methylation: Blocked

Star activity: Large excess of the enzyme results in the appearance of star activity.

Percent Activity in MINOTECH Buffers

L	M	H	SH	A	K
<10	<10	75	50-75	10	100

General reaction mixture:

10U NruI	1μl
10x U _{NruI} or K buffer *	2μl
DNA substrate	<1μg
Sterile ultrapure water	Up to 20 μl

Incubate for 15 min at 37°C

*In the case of U_{NruI} buffer we recommend the addition of BSA to a final concentration of 100 μg/ml.

Frequency of Cutting

λ	Ad-2	Φx174	pUC18	M13mp18	pBR322
5	5	2	0	0	1



Lambda DNA 0.7 % agarose