


Rsa I



5' ...GT▼AC...3'
3' ...CA▲TG...5'

RsaI is a restriction enzyme purified from *Rhodopseudomonas sphaeroides*.

Catalogue No 129-1, 1000 U
 129-2, 3x1000 U

Concentration 10-12u/μl and 40-60u/μl*

*Add an H to cat.# to order the high concentration

Reagents supplied: 10x M and 10x K buffer

Unit substrate: Lambda DNA.

Unit calculation assay conditions: 50 mM NaCl, 10 Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM dithiothreitol, 100 μg/ml BSA. Incubate at 37°C.

Absence of contaminants: 400 units of *Rsa I* do not produce any unspecific cleavage products after 16 hrs incubation with 1 μg of λ DNA at 37°C. After 10-fold overdigestion with *Rsa I*, greater than 95% of the DNA fragments can be ligated and recut with this enzyme.

Storage buffer: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 μg/ml BSA and 50% glycerol. Store at -20°C.

Heat inactivation: 65°C for 20 minutes.

Methylation Sensitivity:

dam methylation: Not sensitive

dcm methylation: Not sensitive

CpG methylation: Blocked by some combinations of overlapping

Note: Cleaves single-stranded DNA slowly.

Percent Activity in MINOTECH Buffers

L	M	H	SH	A	K
75-100	100	50	<10	<10	100


General reaction mixture:

10U RsaI 1μl
10x M or K buffer * 2μl
DNA substrate <1μg
Sterile ultrapure water Up to 20 μl
Incubate for 15 min at 37°C

*In the case of M buffer we recommend the addition of BSA to a final concentration of 100 μg/ml.

Frequency of Cutting

λ	Ad-2	Φx174	pUC18	M13mp18	pBR322
113	83	11	3	19	3



Lambda DNA 1.4 % agarose