


SnaBI



5' ...TAC▼GTA...3'
3' ...ATG▲CAT...5'

SnaBI is a restriction enzyme purified from *Sphaerotilus natans*.

Catalogue No 136-1, 500 U
 136-2, 3x500 U

Concentration 10-12u/μl and 40-60u/μl*

*Add an H to cat.# to order the high concentration

Reagents supplied: 10x U_{SnaBI} and 10x K buffer

Unit substrate: Lambda DNA (EcoRI digest).

Unit calculation assay conditions: 10 mM Bis Tris Propane-HCl (pH 7.0 @ 25°C), 10 mM MgCl₂, 1 mM dithiothreitol, 100 μg/ml BSA. Incubate at 37°C.

Absence of contaminants: 10 units of SnaBI do not produce any unspecific cleavage products after 16 hrs incubation with 1 μg of λ DNA/EcoRI digest at 37°C. After 10-fold overdigestion with SnaBI, greater than 98% of the DNA fragments can be ligated and recut with this enzyme.

Storage buffer: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 μg/ml BSA and 50% glycerol. Store at -20°C.

Heat inactivation: 80°C for 20 minutes.

Methylation Sensitivity:

dam methylation: Not sensitive

dcm methylation: Not sensitive

CpG methylation: Blocked

Percent Activity in MINOTECH Buffers

L	M	H	SH	A	K
50-75	50	25	<10	100	100

General reaction mixture:

10U SnaBI	1μl
10x U _{SnaBI} or K buffer *	2μl
DNA substrate	<1μg
Sterile ultrapure water	Up to 20 μl

Incubate for 15 min at 37°C

*In the case of U_{SnaBI} buffer we recommend the addition of BSA to a final concentration of 100 μg/ml.

Frequency of Cutting

λ	Ad-2	Φx174	pUC18	M13mp18	pBR322
1	0	0	0	1	0



Lambda DNA 0.7 % agarose