


Sty I



5' ...C▼CWWGG...3'
3' ...GGWWC▲C...5'

StyI is a restriction enzyme purified from *E.coli* WA921/pST27 hsd+.

Catalogue No 141-1, 4000 U
141-2, 3x4000 U

Concentration 10-12u/μl and 40-60u/μl*

*Add an H to cat.# to order the high concentration

Reagents supplied: 10x H buffer

Unit substrate: Lambda DNA.

Unit calculation assay conditions: 100 mM NaCl, 50 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM dithiothreitol, 100 μg/ml BSA. Incubate at 37°C.

Absence of contaminants: 50 units of *StyI* do not produce any unspecific cleavage products after 16 hrs incubation with 1 μg of λ DNA at 37°C. After 50-fold overdigestion with *StyI*, greater than 98% of the DNA fragments can be ligated and recut with this enzyme.

Storage buffer: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 200 μg/ml BSA and 50% glycerol. Store at -20°C.

Heat inactivation: 65°C for 20 minutes.

Methylation Sensitivity:

dam methylation: Not sensitive
dcm methylation: Not sensitive
CpG methylation: Not sensitive

Star activity: Conditions of low ionic strength, high enzyme concentration, glycerol concentration >5%, or pH>8.0 may result in star activity

Percent Activity in MINOTECH Buffers

L	M	H	SH	A	K
25-50	75-100	100	75-100	<10	50


General reaction mixture:

10U *StyI* 1μl
10x H buffer * 2μl
DNA substrate <1μg
Sterile ultrapure water Up to 20 μl
Incubate for 15 min at 37°C

*We recommend the addition of BSA to a final concentration of 100 μg/ml.

Frequency of Cutting

λ	Ad-2	Φx174	pUC18	M13mp18	pBR322
10	45	0	0	0	1



Lambda DNA 1.0 % agarose